arXiv:1606.03525v1 [math.PR] 11 Jun 2016

Time varying axially symmetric vector random fields on the sphere

Chunsheng Ma*

Abstract This paper presents a general form of the covariance matrix structure for a vector random field that is axially symmetric and mean square continuous on the sphere and provides a series representation for a longitudinally reversible one. The series representation is somehow an imitator of the covariance matrix function, and both of them have simpler forms than those proposed in the literature in terms of the associated Legendre functions and are useful for modeling and simulation. Also, a general form of the covariance matrix structure is derived for a spatio-temporal vector random field that is axially symmetric and mean square continuous over the sphere, and a series representation is given for a longitudinally reversible one.

Keywords Covariance matrix function · Elliptically contoured random field · Gaussian random field · Isotropic · Longitudinally reversible · Stationary

Mathmatics Subject Classification (2010) 60G60 · 62M10 · 62M30

*Chunsheng Ma Department of Mathematics, Statistics, and Physics, Wichita State University, Wichita, Kansas 67260-0033, USA e-mail: chunsheng.ma@wichita.edu

1 Introduction

Axially symmetric random fields on the three-dimensional sphere were introduced by [16] more than fifty years ago. This kind of spatial or spatio-temporal models looks like reasonable for the practical applications ([4], [14], [17], [18], [19], [36]), but its complicated covariance structure and series expansion ([15], [16], [36]) have made it very hard to fit real data. This calls for a more detailed investigation, and motivates us here to search for simple forms of the correlation structure and series expansion for a scalar, vector, or time varying random field that is axially symmetric and mean square continuous on the sphere. The established theories differ from, but are not more complicated than, those of the scalar, vector, or time varying random field that is isotropic and mean square continuous on spheres ([2], [3], [5], [6], [7], [8], [10], [11], [13], [21], [22], [24] - [30], [32], [33], [34], [39], [40], [41]).

Denote by $\mathbb{S}^2$ the spherical shell of radius 1 and center $\mathbf{0}$ in $\mathbb{R}^3$, i.e., $\mathbb{S}^2 = \{\|\mathbf{x}\| = 1, \mathbf{x} \in \mathbb{R}^3\}$, where $\|\mathbf{x}\|$ is the Euclidean norm of $\mathbf{x} \in \mathbb{R}^3$. Using spherical coordinates, a point $\mathbf{x} \in \mathbb{S}^2$ is determined by the longitude θ and the latitude φ, and is designated as $\mathbf{x} = (\varphi, \theta)$, where $0 \leq \theta \leq 2\pi$ and $0 \leq \varphi \leq \pi$.

An m-variate real and second-order random field $\{\mathbf{Z}(\mathbf{x}), \mathbf{x} = (\varphi, \theta) \in \mathbb{S}^2\}$ is said to be axially symmetric, if it mean function $\mathrm{E}\mathbf{Z}(\mathbf{x})$ depends only on φ, and its covariance matrix function $\mathrm{cov}(\mathbf{Z}(\mathbf{x}_1), \mathbf{Z}(\mathbf{x}_2))$ depends on φ_1, φ_2, and $\theta_1 - \theta_2$. The covariance matrix function is denoted by $\mathbf{C}(\varphi_1, \varphi_2, \theta)$, or

$$\mathbf{C}(\varphi_1, \varphi_2, \theta_1 - \theta_2) = \mathrm{E}\{(\mathbf{Z}(\mathbf{x}_1) - \mathrm{E}\mathbf{Z}(\mathbf{x}_1))(\mathbf{Z}(\mathbf{x}_2) - \mathrm{E}\mathbf{Z}(\mathbf{x}_2))'\}, \quad \mathbf{x}_1 = (\varphi_1, \theta_1), \mathbf{x}_2 = (\varphi_2, \theta_2) \in \mathbb{S}^2.$$

Moreover, the random field or its covariance matrix function is said to be longitudinally reversible [36], if

$$\mathbf{C}(\varphi_1, \varphi_2, -\theta) = \mathbf{C}(\varphi_1, \varphi_2, \theta), \qquad \varphi_1, \varphi_2 \in [0, \pi], \ \theta \in [-2\pi, 2\pi].$$

In a spatio-temporal setting, an m-variate real and second-order random field $\{\mathbf{Z}(\mathbf{x}; t), \mathbf{x} \in \mathbb{S}^2, t \in \mathbb{T}\}$ is said to be axially symmetric on $\mathbb{S}^2$ and stationary over the temporal domain $\mathbb{T}$, if it mean function $\mathrm{E}\mathbf{Z}(\mathbf{x}; t)$ depends only on φ, and its covariance matrix function $\mathrm{cov}(\mathbf{Z}(\mathbf{x}_1; t_1), \mathbf{Z}(\mathbf{x}_2; t_2))$ depends on φ_1, φ_2, $\theta_1 - \theta_2$, and $t_1 - t_2$, where $\mathbb{T} = \mathbb{R}$ or $\mathbb{T}$. The latter is denoted by $\mathbf{C}(\varphi_1, \varphi_2, \theta; t)$,

or

$$\mathbf{C}(\varphi_1,\varphi_2,\theta_1-\theta_2;t_1-t_2)=\mathrm{E}\{(\mathbf{Z}(\mathbf{x}_1;t_1)-\mathrm{E}\mathbf{Z}(\mathbf{x}_1;t_1))(\mathbf{Z}(\mathbf{x}_2;t_2)-\mathrm{E}\mathbf{Z}(\mathbf{x}_2;t_2))'\},$$

$$\mathbf{x}_k=(\varphi_k,\theta_k)\in\mathbb{S}^2,\ t_k\in\mathbb{T},\ k=1,2.$$

Two fundamental properties of this $m\times m$ matrix function are: (i) $\mathbf{C}(\varphi_1,\varphi_2,\theta;t)=(\mathbf{C}(\varphi_2,\varphi_1,-\theta;-t))'$, and (ii) inequality

$$\sum_{i=1}^{n}\sum_{j=1}^{n}\mathbf{a}_i'\mathbf{C}(\varphi_i,\varphi_j,\theta_i-\theta_j;t_i-t_j)\mathbf{a}_j\geq 0 \tag{1}$$

holds for every $n\in\mathbb{N}$, any $\varphi_i\in[0,\pi],\theta_i\in[0,2\pi]$, $t_i\in\mathbb{T}$, and $\mathbf{a}_i\in\mathbb{R}^m$ $(i=1,2,\ldots,n)$, where $\mathbb{N}$ stands for the set of positive integers. On the other hand, given an $m\times m$ matrix function with these properties, there exists an m-variate Gaussian or elliptically contoured random field $\{\mathbf{Z}(\mathbf{x};t),\mathbf{x}\in\mathbb{S}^2,t\in\mathbb{T}\}$ with $\mathbf{C}(\varphi_1,\varphi_2,\theta;t)$ as its covariance matrix function [23].

Section 2 derives a general form of the covariance matrix structure of an m-variate axially symmetric and mean square continuous random field on $\mathbb{S}^2$ and a series representation of a longitudinally reversible one, starting from a close look at the scalar case of [16]. Moving successively, a general covariance matrix form is given for an m-vatiate spatio-temporal random field that is axially symmetric and mean square continuous on $\mathbb{S}^2$ and stationary on $\mathbb{T}$, and a series representation is offered to a longitudinally reversible one over the $\mathbb{S}^2$. Theorems 1-3 and 5-7 are proved in Section 4.

2 Covariance matrix structures and series representations

Suppose that $\{\mathbf{Z}(\mathbf{x}),\mathbf{x}\in\mathbb{S}^2\}$ is an m-variate axially symmetric and mean square continuous random field. This section derives a general form of its covariance matrix function, which does not look as complicated as that in [16] does, and presents a series representation for a longitudinally reversible one, which is useful for modeling and simulation. An isotropic version is treated in [24], [26].

To begin with, let us take a look at (12) of [16] for the covariance function of a scalar axially

symmetric random field $\{Z(\varphi,\theta),\varphi\in[0,\pi],\theta\in[0,2\pi]\}$, which reads (in our notions)

$$C(\varphi_1,\varphi_2,\theta)=\sum_{n=0}^{\infty}(b_n(\varphi_1,\varphi_2)\cos(n\theta)+a_n(\varphi_1,\varphi_2)\sin(n\theta)),\quad \varphi_1,\varphi_2\in[0,\pi],\theta\in[-2\pi,2\pi],\quad (2)$$

where

$$\begin{aligned} b_n(\varphi_1,\varphi_2) &= \sum_{i=n}^{\infty}\sum_{j=n}^{\infty} b_{n,ij}P_i^n(\cos\varphi_1)P_j^n(\cos\varphi_2),\\ a_n(\varphi_1,\varphi_2) &= \sum_{i=n}^{\infty}\sum_{j=n}^{\infty} a_{n,ij}P_i^n(\cos\varphi_1)P_j^n(\cos\varphi_2),\qquad n\in\mathbb{N}_0, \end{aligned}$$

$b_{n,ij}$ and $a_{n,ij}$ are certain constants, $P_k^n(x)$ are the associated Legendre function [1], and $\mathbb{N}_0$ denotes the set of nonnegative integers. Simply speaking, (2) is a Fourier series expansion of $C(\varphi_1,\varphi_2,\theta)$ in terms of $\theta\in[-2\pi,2\pi]$, but using or not the associated Legendre function is not so important there, as is indicated by Theorem 1.

Theorem 1 *For an axially symmetric and mean square continuous random field $\{Z(\mathbf{x}),\mathbf{x}\in\mathbb{S}^2\}$, its covariance function $C(\varphi_1,\varphi_2,\theta)$ is of the form*

$$C(\varphi_1,\varphi_2,\theta)=\sum_{n=0}^{\infty}\{b_n(\varphi_1,\varphi_2)\cos(n\theta)+a_n(\varphi_1,\varphi_2)\sin(n\theta)\},\quad \varphi_1,\varphi_2\in[0,\pi],\ \theta\in[-2\pi,2\pi],\ (3)$$

where $b_n(\varphi_1,\varphi_2)$ is a covariance function on $[0,\pi]$ for each fixed $n\in\mathbb{N}_0$, and $\sum_{n=0}^{\infty} b_n(\varphi_1,\varphi_2)$ converges.

Although $b_n(\varphi_1,\varphi_2)$ has to be a covariance function for each $n\in\mathbb{N}_0$, it is not clear how to interpret $a_n(\varphi_1,\varphi_2)$. No matter what $a_n(\varphi_1,\varphi_2)$ looks like, it must make (3) a positive definite function. An example of (3) is illustrated in Example 1, where $a_n(\varphi_1,\varphi_2)$ is either non-negatively or negatively proportional to $b_n(\varphi_1,\varphi_2)$.

Example 1 For a constant λ with $|\lambda|\le 1$,

$$C(\varphi_1,\varphi_2,\theta)=\sum_{n=0}^{\infty} b_n(\varphi_1,\varphi_2)(\cos(n\theta)+\lambda\sin(n\theta)),\quad \varphi_1,\varphi_2\in[0,\pi],\ \theta\in[-2\pi,2\pi],$$

is the covariance function of an axially symmetric Gaussian or elliptically contoured random field on $\mathbb{S}^2$, where $b_n(\varphi_1, \varphi_2)$ is a covariance function on $[0, \pi]$ for each fixed $n \in \mathbb{N}_0$, and $\sum_{n=0}^{\infty} b_n(\varphi_1, \varphi_2)$ converges. This follows from Theorem 8 of [23], noticing that $\cos(n\theta) + \lambda \sin(n\theta)$ is a positive definite function of $\theta \in [-2\pi, 2\pi]$ for each $n \in \mathbb{N}_0$. Of course, many $b_n(\varphi_1, \varphi_2)$ are available for selection.

Corollary 1.1 *A longitudinally reversible covariance function $C(\varphi_1, \varphi_2, \theta)$ takes the form*

$$C(\varphi_1, \varphi_2, \theta) = \sum_{n=0}^{\infty} b_n(\varphi_1, \varphi_2) \cos(n\theta), \quad \varphi_1, \varphi_2 \in [0, \pi], \ \theta \in [-2\pi, 2\pi]. \tag{4}$$

Conversely, given a function $C(\varphi_1, \varphi_2, \theta)$ of the form (4) with summable $\sum_{n=0}^{\infty} b_n(\varphi_1, \varphi_2)$ and each $b_n(\varphi_1, \varphi_2)$ being a covariance function on $[0, \pi]$, there exists a longitudinally reversible Gaussian or elliptical contoured random field on $\mathbb{S}^2$ with it as the covariance function.

In fact, (4) follows from (3) and $C(\varphi_1, \varphi_2, \theta) = C(\varphi_1, \varphi_2, -\theta)$; see also Proposition 3 of [15]. One way to establish the second part of Corollary 1.1 is to verify the positive definiteness of (4) and to use Theorem 8 of [23]. Instead, we give a series representation for a random field with (4) as its covariance function, which is useful for modeling and simulation.

Theorem 2. *Assume that stochastic processes $\{V_{n1}(\varphi), \varphi \in [0, \pi]\}$ and $\{V_{n2}(\varphi), \varphi \in [0, \pi]\}$ have mean 0 and covariance function $b_n(\varphi_1, \varphi_2)$ for each $n \in \mathbb{N}_0$, and that for all $n \in \mathbb{N}_0$, $\{V_{n1}(\varphi), \varphi \in [0, \pi]\}$ and $\{V_{n2}(\varphi), \varphi \in [0, \pi]\}$ are independent. If $\sum_{n=0}^{\infty} b_n(\varphi_1, \varphi_2)$ converges, then*

$$Z(\varphi, \theta) = \sum_{n=0}^{\infty} (V_{n1}(\varphi) \cos(n\theta) + V_{n2}(\varphi) \sin(n\theta)), \qquad \varphi \in [0, \pi], \ \theta \in [0, 2\pi], \tag{5}$$

is a longitudinally reversible random field on $\mathbb{S}^2$ with mean 0 and covariance function (4).

Somehow (5) is an imitator of (3) but not (4). It would be of interest to generate an axially symmetric random field on $\mathbb{S}^2$. The vector case is slightly more complicated than the scalar case, since a covariance matrix function is not necessarily symmetric.

Theorem 3 *If an m-variate mean square continuous random field $\{\mathbf{Z}(\mathbf{x}), \mathbf{x} \in \mathbb{S}^2\}$ is axially symmetric, then $\frac{\mathbf{C}(\varphi_1,\varphi_2,\theta)+(\mathbf{C}(\varphi_1,\varphi_2,\theta))'}{2}$ is of the form*

$$\frac{\mathbf{C}(\varphi_1,\varphi_2,\theta)+(\mathbf{C}(\varphi_1,\varphi_2,\theta))'}{2} = \sum_{n=0}^{\infty} \{\mathbf{B}_n(\varphi_1,\varphi_2)\cos(n\theta) + \mathbf{A}_n(\varphi_1,\varphi_2)\sin(n\theta)\}, \quad \varphi_1,\varphi_2 \in [0,\pi],\ \theta \in [-2\pi, 2\pi], \tag{6}$$

where $\mathbf{B}_n(\varphi_1,\varphi_2)$ ($n \in \mathbb{N}_0$) are $m \times m$ symmetric matrices, $\sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2)$ converges, and, for each fixed $n \in \mathbb{N}_0$, $\mathbf{B}_n(\varphi_1,\varphi_2)$ is a covariance matrix function on $[0,\pi]$.

Example 2 For a constant λ with $|\lambda| \leq 1$,

$$\mathbf{C}(\varphi_1,\varphi_2,\theta) = \sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2)(\cos(n\theta) + \lambda\sin(n\theta)), \quad \varphi_1,\varphi_2 \in [0,\pi],\ \theta \in [-2\pi,2\pi],$$

is the covariance matrix function of an m-variate axially symmetric Gaussian or elliptically contoured random field on $\mathbb{S}^2$, if $\mathbf{B}_n(\varphi_1,\varphi_2)$ is an $m \times m$ covariance matrix function on $[0,\pi]$ for each fixed $n \in \mathbb{N}_0$, and $\sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2)$ converges.

Corolloary 3.1 *If $\mathbf{C}(\varphi_1,\varphi_2,\theta)$ is longitudinally reversible, then*

$$\frac{\mathbf{C}(\varphi_1,\varphi_2,\theta) + (\mathbf{C}(\varphi_1,\varphi_2,\theta))'}{2} = \sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2)\cos(n\theta), \quad \varphi_1,\varphi_2 \in [0,\pi],\ \theta \in [-2\pi,2\pi]. \tag{7}$$

Corollary 3.2 *An $m \times m$ matrix function*

$$\mathbf{C}(\varphi_1,\varphi_2,\theta) = \sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2)\cos(n\theta), \quad \varphi_1,\varphi_2 \in [0,\pi],\ \theta \in [-2\pi,2\pi], \tag{8}$$

is the covariance matrix function of an m-variate Gaussian or elliptically contoured random field on $\mathbb{S}^2$ if and only if $\mathbf{B}_n(\varphi_1,\varphi_2)$ is an $m \times m$ covariance matrix function on $[0,\pi]$ for each fixed $n \in \mathbb{N}_0$, and $\sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2)$ converges.

The matrix $\mathbf{B}_n(\varphi_1,\varphi_2)$ in (7) has to be symmetric in the sense that $\mathbf{B}_n(\varphi_1,\varphi_2) = (\mathbf{B}_n(\varphi_1,\varphi_2))'$. But, it is not necessarily so in (8) or the next theorem, whose proof is similar to that of Theorem 2 and is thus omitted.

Theorem 4 *Assume that m-variate stochastic processes $\{\mathbf{V}_{n1}(\varphi), \varphi \in [0,\pi]\}$ and $\{\mathbf{V}_{n2}(\varphi), \varphi \in [0,\pi]\}$ have mean $\mathbf{0}$ and covariance matrix function $\mathbf{B}_n(\varphi_1,\varphi_2)$ for each $n \in \mathbb{N}_0$, and that for all $n \in \mathbb{N}_0$, $\{\mathbf{V}_{n1}(\varphi), \varphi \in [0,\pi]\}$ and $\{\mathbf{V}_{n2}(\varphi), \varphi \in [0,\pi]\}$ are independent. If $\sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2)$ converges, then*

$$\mathbf{Z}(\varphi,\theta) = \sum_{n=0}^{\infty} (\mathbf{V}_{n1}(\varphi)\cos(n\theta) + \mathbf{V}_{n2}(\varphi)\sin(n\theta)), \qquad \varphi \in [0,\pi],\ \theta \in [0,2\pi], \tag{9}$$

is an m-variate longitudinally reversible random field on $\mathbb{S}^2$ with mean $\mathbf{0}$ and covariance matrix function (8).

Example 3 Let

$$\mathbf{B}_{n,ij}(\varphi_1,\varphi_2) = \begin{cases} \frac{1}{(b_i(\varphi_1)+b_j(\varphi_2))\pi}, & n = 0, \\ \frac{2}{(n^2+b_i(\varphi_1)+b_j(\varphi_2))\pi}, & n \in \mathbb{N},\ \varphi_1,\varphi_2 \in [0,\pi],\quad i,j = 1,\ldots,m, \end{cases}$$

where $b_i(\varphi)$ $(i = 1,\ldots,m)$ are positive functions on $[0,\pi]$. It can be verified that each $\mathbf{B}_n(\varphi_1,\varphi_2)$ is a covariance matrix function on $[0,\pi]$. The direct/cross covariance functions of (9) are

$$\begin{aligned} C_{ij}(\varphi_1,\varphi_2,\theta) &= \frac{\cosh\left((\pi - |\theta|)\sqrt{b_i(\varphi_1)+b_j(\varphi_2)}\right)}{\sqrt{b_i(\varphi_1)+b_j(\varphi_2)}\sinh(\sqrt{b_i(\varphi_1)+b_j(\varphi_2)}\pi)}, \\ &\qquad \varphi_1,\varphi_2 \in [0,\pi],\ \theta \in [-2\pi,2\pi],\ i,j = 1,\ldots,m, \end{aligned}$$

based on the identity (see, e.g., page 577 of [38])

$$\sum_{n=1}^{\infty} \frac{\cos(n\theta)}{n^2+a^2} = \frac{\pi\cosh((\pi-\theta)a)}{2a\sinh(\pi a)} - \frac{1}{2a^2}, \quad a > 0,\ \theta \in [0,2\pi].$$

3 Time varying axially symmetric vector random fields

This section deals with the covariance matrix structure of an m-variate random field $\{\mathbf{Z}(\mathbf{x};t), \mathbf{x} \in \mathbb{S}^2, t \in \mathbb{T}\}$ that is axially symmetric and mean square continuous on $\mathbb{S}^2$ and stationary on $\mathbb{T}$, and presents a series representation for a longitudinally reversible one. An isotropic case is studied

in [28]. The connection of the Gaussianity between the random field and its coefficients in the series representation is explored in Theorem 6. Some arguments similar to the purely spatial case are not repeated.

Theorem 5 *If an m-variate random field $\{\mathbf{Z}(\mathbf{x};t), \mathbf{x} \in \mathbb{S}^2, t \in \mathbb{T}\}$ is axially symmetric and mean square continuous on $\mathbb{S}^2$ and stationary on $\mathbb{T}$, then*

$$\begin{aligned}&\tfrac{\mathbf{C}(\varphi_1,\varphi_2,\theta;t)+(\mathbf{C}(\varphi_1,\varphi_2,\theta;t))'+\mathbf{C}(\varphi_1,\varphi_2,\theta;-t)+(\mathbf{C}(\varphi_1,\varphi_2,\theta;-t))'}{2}\\ &= \sum_{n=0}^{\infty} \left\{\mathbf{B}_n(\varphi_1,\varphi_2;t)\cos(n\theta) + \mathbf{A}_n(\varphi_1,\varphi_2;t)\sin(n\theta)\right\}, \qquad (10)\\ &\qquad\qquad \varphi_1,\varphi_2 \in [0,\pi],\ \theta \in [-2\pi,2\pi], t \in \mathbb{T},\end{aligned}$$

where $\mathbf{B}_n(\varphi_1,\varphi_2;t)$ $(n \in \mathbb{N}_0)$ are $m \times m$ symmetric matrices and $\sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2;t)$ converges, and, for each fixed $n \in \mathbb{N}_0$, $\mathbf{B}_n(\varphi_1,\varphi_2;t)$ is a covariance matrix function on $[0,\pi]\times\mathbb{T}$.

Corollary 5.1 *If $\{\mathbf{Z}(\mathbf{x};t), \mathbf{x} \in \mathbb{S}^2, t \in \mathbb{T}\}$ is longitudinally reversible on $\mathbb{S}^2$, then*

$$\begin{aligned}&\tfrac{\mathbf{C}(\varphi_1,\varphi_2,\theta;t)+(\mathbf{C}(\varphi_1,\varphi_2,\theta;t))'+\mathbf{C}(\varphi_1,\varphi_2,\theta;-t)+(\mathbf{C}(\varphi_1,\varphi_2,\theta;-t))'}{2} = \sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2;t)\cos(n\theta),\\ &\qquad\qquad \varphi_1,\varphi_2 \in [0,\pi],\ \theta \in [-2\pi,2\pi], t \in \mathbb{T}.\end{aligned} \qquad (11)$$

Corollary 5.2 *An $m \times m$ matrix function*

$$\mathbf{C}(\varphi_1,\varphi_2,\theta;t) = \sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2;t)\cos(n\theta), \quad \varphi_1,\varphi_2 \in [0,\pi],\ \theta \in [-2\pi,2\pi],\ t \in \mathbb{T}, \qquad (12)$$

is the covariance matrix function of an m-variate random field on $\mathbb{S}^2 \times \mathbb{T}$ that is longitudinally reversible on $\mathbb{S}^2$ and stationary on $\mathbb{T}$ if and only if $\sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1,\varphi_2;t)$ converges, and, for each fixed $n \in \mathbb{N}_0$, $\mathbf{B}_n(\varphi_1,\varphi_2;t)$ is a covariance matrix function on $[0,\pi]\times\mathbb{T}$.

Theorem 6 *Assume that m-variate stochastic processes $\{\mathbf{V}_{n1}(\varphi;t), \varphi \in [0,\pi], t \in \mathbb{T}\}$ and $\{\mathbf{V}_{n2}(\varphi;t), \varphi \in [0,\pi], t \in \mathbb{T}\}$ have mean $\mathbf{0}$ and covariance matrix function $\mathbf{B}_n(\varphi_1,\varphi_2;t)$ for each $n \in \mathbb{N}_0$, and that for all $n \in \mathbb{N}_0$, $\{\mathbf{V}_{n1}(\varphi;t), \varphi \in [0,\pi], t \in \mathbb{T}\}$ and $\{\mathbf{V}_{n2}(\varphi;t), \varphi \in [0,\pi], t \in \mathbb{T}\}$*

are independent. If $\sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1, \varphi_2; t)$ *converges, then*

$$\mathbf{Z}(\varphi, \theta; t) = \sum_{n=0}^{\infty} (\mathbf{V}_{n1}(\varphi; t) \cos(n\theta) + \mathbf{V}_{n2}(\varphi; t) \sin(n\theta)), \qquad \varphi \in [0, \pi],\ \theta \in [0, 2\pi], t \in \mathbb{T}, \quad (13)$$

is an m*-variate random field longitudinally reversible on* $\mathbb{S}^2$ *and stationary on* $\mathbb{T}$*, with mean* $\mathbf{0}$ *and covariance matrix function*

$$\mathbf{C}(\varphi_1, \varphi_2, \theta; t) = \sum_{n=0}^{\infty} \mathbf{B}_n(\varphi_1, \varphi_2; t) \cos(n\theta), \quad \varphi_1, \varphi_2 \in [0, \pi],\ \theta \in [-2\pi, 2\pi],\ t \in \mathbb{T}.$$

Moreover, $\mathbf{Z}(\varphi, \theta; t)$ *is Gaussian if and only if all* $\mathbf{V}_{n1}(\varphi; t)$ *and* $\mathbf{V}_{n2}(\varphi; t)$ *(*$n \in \mathbb{N}_0$*) are Gaussian.*

Example 4 Let $\mathbf{B}(\varphi_1, \varphi_2; t)$ be an $m \times m$ covariance matrix function on $[0, \pi] \times \mathbb{T}$ and all its entries be less than 1 in absolute value.

(i) In Theorem 6 choose

$$\mathbf{B}_n(\varphi_1, \varphi_2; t) = \begin{cases} \frac{2}{n} \left(\mathbf{B}(\varphi_1, \varphi_2; t)\right)^{\circ n}, & n \in \mathbb{N}, \\ \mathbf{0}, & n = 0,\ \varphi_1, \varphi_2 \in [0, \pi],\ t \in \mathbb{T}, \end{cases}$$

where $\mathbf{B}^{\circ p}$ denotes the Hadamard p power of $\mathbf{B} = (b_{ij})$, whose entries are b_{ij}^p, the p power of $b_{ij}, i, j = 1, \ldots, m$. The direct/cross covariance functions of (13) are

$$C_{ij}(\varphi_1, \varphi_2, \theta; t) = -\ln(1 - 2b_{ij}(\varphi_1, \varphi_2; t) \cos\theta + b_{ij}^2(\varphi_1, \varphi_2; t)),$$

$$\varphi_1, \varphi_2 \in [0, \pi],\ \theta \in [-2\pi, 2\pi],\ t \in \mathbb{T}, i, j = 1, \ldots, m,$$

in view of the identity

$$\sum_{n=1}^{\infty} \frac{a^n}{n} \cos(n\theta) = -\frac{1}{2} \ln(1 - 2a \cos\theta + a^2), \qquad |a| < 1.$$

(ii) Choose

$$\mathbf{B}_n(\varphi_1, \varphi_2) = \begin{cases} \left(\mathbf{B}(\varphi_1, \varphi_2; t)\right)^{\circ n}, & n \in \mathbb{N}, \\ \mathbf{1}, & n = 0,\ \varphi_1, \varphi_2 \in [0, \pi],\ t \in \mathbb{T}, \end{cases}$$

where $\mathbf{1}$ is an $m \times m$ matrix with all entries 1. Then the direct/cross covariance functions of (13) are

$$C_{ij}(\varphi_1, \varphi_2, \theta; t) = \frac{1 - b_{ij}^2(\varphi_1, \varphi_2; t)}{1 - 2b_{ij}(\varphi_1, \varphi_2; t)\cos\theta + b_{ij}^2(\varphi_1, \varphi_2; t)},$$
$$\varphi_1, \varphi_2 \in [0, \pi],\ \theta \in [-2\pi, 2\pi], t \in \mathbb{T},\ i, j = 1, \ldots, m,$$

which follows from the identity

$$\sum_{n=0}^{\infty} a^n \cos(n\theta) = \frac{1 - a^2}{1 - 2a\cos\theta + a^2}, \qquad |a| < 1.$$

A sufficient condition is given in the following theorem for an $m \times m$ matrix function to be the covariance matrix function of an m-variate Gaussian or elliptically contoured random field on $\mathbb{S}^2 \times \mathbb{T}$ that is axially symmetric and mean square continuous on $\mathbb{S}^2$ and stationary on $\mathbb{T}$.

Theorem 7. *For each $n \in \mathbb{N}_0$, if the inequality*

$$\sum_{i=1}^{l}\sum_{j=1}^{l}(\mathbf{u}_i'\mathbf{B}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{u}_j + \mathbf{v}_i'\mathbf{B}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{v}_j + \mathbf{u}_i'\mathbf{A}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{v}_j - \mathbf{v}_i'\mathbf{A}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{u}_j) \geq 0 \tag{14}$$

holds for every $l \in \mathbb{N}$, any $\mathbf{u}_i \in \mathbb{R}^m, \mathbf{v}_i \in \mathbb{R}^m$, $t_i \in \mathbb{T}$, and $\varphi_i \in [0, \pi]$, then there exists an m-variate Gaussian or elliptically contoured random field on $\mathbb{S}^2 \times \mathbb{T}$ with the covariance matrix function

$$\mathbf{C}(\varphi_1, \varphi_2, \theta; t) = \sum_{n=0}^{\infty} \{\mathbf{B}_n(\varphi_1, \varphi_2; t)\cos(n\theta) + \mathbf{A}_n(\varphi_1, \varphi_2; t)\sin(n\theta)\},$$
$$\varphi_1, \varphi_2 \in [0, \pi],\ \theta \in [-2\pi, 2\pi], t \in \mathbb{T}. \tag{15}$$

Inequality (14) ensures that each level n of (15), $\mathbf{B}_n(\varphi_1, \varphi_2; t)\cos(n\theta) + \mathbf{A}_n(\varphi_1, \varphi_2; t)\sin(n\theta)$, satisfies inequality (1). It would be of interest to see whether (14) is also a necessary condition.

4 Proofs

4.1 Proof of Theorem 1

Recall that the mean square continuity of $\{Z(\mathbf{x}), \mathbf{x} \in \mathbb{S}^2\}$ means

$$\mathrm{E}|Z(\varphi_1, \theta_1) - Z(\varphi_2, \theta_2)|^2 \to 0, \quad \text{as } \varphi_1 \to \varphi_2, \theta_1 \to \theta_2.$$

This implies that $C(\varphi_1, \varphi_2, \theta)$ is continuous with respect to $\theta \in [-2\pi, 2\pi]$ for fixed $\varphi_1, \varphi_2 \in [0, \pi]$. Indeed, it follows from the Cauchy-Schwartz inequality that, for $\theta_1, \theta_2 \geq 0$,

$$\begin{aligned}
& |C(\varphi_1, \varphi_2, \theta_1) - C(\varphi_1, \varphi_2, \theta_2)| \\
= \; & |\mathrm{E}Z(\varphi_1, \theta_1)(Z(\varphi_2, 0) - \mathrm{E}Z(\varphi_2, 0)) - \mathrm{E}Z(\varphi_1, \theta_2)(Z(\varphi_2, 0) - \mathrm{E}Z(\varphi_2, 0))| \\
= \; & |\mathrm{E}\{(Z(\varphi_1, \theta_1) - Z(\varphi_1, \theta_2))(Z(\varphi_2, 0) - \mathrm{E}Z(\varphi_2, 0))\}| \\
\leq \; & \mathrm{var}(Z(\varphi_2))\mathrm{E}|Z(\varphi_1, \theta_1) - Z(\varphi_1, \theta_2)|^2 \\
\to \; & 0, \qquad \theta_1 \to \theta_2,
\end{aligned}$$

and, for $\theta_1, \theta_2 \leq 0$,

$$\begin{aligned}
& |C(\varphi_1, \varphi_2, \theta_1) - C(\varphi_1, \varphi_2, \theta_2)| \\
= \; & |\mathrm{E}(Z(\varphi_1, 0) - \mathrm{E}Z(\varphi_1, 0))Z(\varphi_2, -\theta_1) - \mathrm{E}(Z(\varphi_1, 0) - \mathrm{E}Z(\varphi_1, 0))Z(\varphi_2, -\theta_2)| \\
= \; & |\mathrm{E}\{(Z(\varphi_1, 0) - \mathrm{E}Z(\varphi_1, 0))(Z(\varphi_2, -\theta_1) - Z(\varphi_2, -\theta_2))\}| \\
\leq \; & \mathrm{var}(Z(\varphi_1))\mathrm{E}|Z(\varphi_2, -\theta_1) - Z(\varphi_2, -\theta_2)|^2 \\
\to \; & 0, \qquad \theta_1 \to \theta_2.
\end{aligned}$$

Thus, as a continuous function of $\theta \in [-2\pi, 2\pi]$, $C(\varphi_1, \varphi_2, \theta)$ takes the Fourier series expansion (3), since $\{\cos(n\theta), \sin(n\theta), n \in \mathbb{N}_0\}$ consists of an orthogonal basis of $L^2[-2\pi, 2\pi]$. The convergence of $\sum_{n=0}^{\infty} b_n(\varphi_1, \varphi_2)$ is due to the existence of $C(\varphi_1, \varphi_2, 0)$.

What remains is to confirm that each coefficient in (3), $b_n(\varphi_1, \varphi_2)$, is a covariance function. For this purpose, consider a stochastic process

$$W_n(\varphi) = \int_0^{2\pi} Z(\varphi, \theta)\cos(n\theta)d\theta, \qquad \varphi \in [0, \pi],$$

for each fixed $n \in \mathbb{N}_0$. In the light of (3), the covariance function of $\{W_n(\varphi), \varphi \in [0,\pi]\}$ is obtained as follows,

$$
\begin{aligned}
& \operatorname{cov}(W_n(\varphi_1), W_n(\varphi_2)) \\
= & \int_0^{2\pi}\int_0^{2\pi} \operatorname{cov}(Z(\varphi_1,\theta_1), Z(\varphi_2,\theta_2))\cos(n\theta_1)\cos(n\theta_2)d\theta_1 d\theta_2 \\
= & \int_0^{2\pi}\int_0^{2\pi} C(\varphi_1,\varphi_2,\theta_1-\theta_2)\cos(n\theta_1)\cos(n\theta_2)d\theta_1 d\theta_2 \\
= & \int_0^{2\pi}\int_0^{2\pi} \sum_{k=0}^{\infty}\{b_k(\varphi_1,\varphi_2)\cos(k(\theta_1-\theta_2)) + a_k(\varphi_1,\varphi_2,\theta_1-\theta_2)\sin(k(\theta_1-\theta_2))\}\cos(n\theta_1)\cos(n\theta_2)d\theta_1 d\theta_2 \\
= & \sum_{k=0}^{\infty} b_k(\varphi_1,\varphi_2)\int_0^{2\pi}\int_0^{2\pi}(\cos(k\theta_1)\cos(k\theta_2)+\sin(k\theta_1)\sin(k\theta_2))\cos(n\theta_1)\cos(n\theta_2)d\theta_1 d\theta_2 \\
& + \sum_{k=0}^{\infty} a_k(\varphi_1,\varphi_2)\int_0^{2\pi}\int_0^{2\pi}(\sin(k\theta_1)\cos(k\theta_2)-\cos(k\theta_1)\sin(k\theta_2))\cos(n\theta_1)\cos(n\theta_2)d\theta_1 d\theta_2 \\
= & \sum_{k=0}^{\infty} b_k(\varphi_1,\varphi_2)\left(\int_0^{2\pi}\cos(k\theta_1)\cos(n\theta_1)d\theta_1\int_0^{2\pi}\cos(k\theta_2)\cos(n\theta_2)d\theta_2\right. \\
& \left. + \int_0^{2\pi}\sin(k\theta_1)\cos(n\theta_1)d\theta_1\int_0^{2\pi}\sin(k\theta_2)\cos(n\theta_2)d\theta_2\right) \\
= & \begin{cases} 4\pi^2 b_0(\varphi_1,\varphi_2), & n=0, \\ \pi^2 b_n(\varphi_1,\varphi_2), & n\in\mathbb{N}, \end{cases}
\end{aligned}
$$

which implies that $b_n(\varphi_1,\varphi_2)$ is a covariance function for each $n \in \mathbb{N}_0$.

4.2 Proof of Theorem 2

The right-hand series of (5) is mean square convergent under the convergent assumption of $\sum_{n=0}^{\infty} b_n(\varphi_1,\varphi_2)$, since

$$
\begin{aligned}
& \mathrm{E}\left|\sum_{n=n_1}^{n_1+n_2}(V_{n1}(\varphi)\cos(n\theta)+V_{n2}(\varphi)\sin(n\theta))\right|^2 \\
= & \sum_{n=n_1}^{n_1+n_2}(EV_{n1}^2(\varphi)+EV_{n2}^2(\varphi)) \\
= & 2\sum_{n=n_1}^{n_1+n_2} b_n(\varphi,\varphi)
\end{aligned}
$$

$$\to \quad 0, \qquad n_1, n_2 \to \infty.$$

It is easy to verify that its covariance function is given by (7).

4.3 Proof of Theorem 3

Since $\{\mathbf{Z}(\mathbf{x}), \mathbf{x} \in \mathbb{S}^2\}$ is an m-variate axially symmetric and mean square continuous random field, two scalar random fields $\{Z_i(\varphi_1, \theta) + Z_j(\varphi_2, \theta), \theta \in [0, 2\pi]\}$ and $\{Z_i(\varphi_1, \theta) - Z_j(\varphi_2, \theta), \theta \in [0, 2\pi]\}$ are also axially symmetric and mean square continuous for fixed φ_k $(k = 1, 2)$ and $i, j \in \{1, \ldots, m\}$, with covariance functions

$$\begin{aligned}
& \operatorname{cov}(Z_i(\varphi_1, \theta_1) + Z_j(\varphi_1, \theta_1), Z_i(\varphi_2, \theta_2) + Z_j(\varphi_2, \theta_2)) \\
= \; & C_{ii}(\varphi_1, \varphi_2, \theta_1 - \theta_2) + C_{ij}(\varphi_1, \varphi_2, \theta_1 - \theta_2) + C_{ji}(\varphi_1, \varphi_2, \theta_1 - \theta_2) + C_{jj}(\varphi_1, \varphi_2, \theta_1 - \theta_2), \\
& \operatorname{cov}(Z_i(\varphi_1, \theta_1) - Z_j(\varphi_1, \theta_1), Z_i(\varphi_2, \theta_2) - Z_j(\varphi_2, \theta_2)) \\
= \; & C_{ii}(\varphi_1, \varphi_2, \theta_1 - \theta_2) - C_{ij}(\varphi_1, \varphi_2, \theta_1 - \theta_2) - C_{ji}(\varphi_1, \varphi_2, \theta_1 - \theta_2) + C_{jj}(\varphi_1, \varphi_2, \theta_1 - \theta_2),
\end{aligned}$$

$$\varphi_1, \varphi_2, \theta_1, \theta_2 \in [0, 2\pi],$$

respectively. It follows from Theorem 1 that

$$\begin{aligned}
& C_{ii}(\varphi_1, \varphi_2, \theta) + C_{ij}(\varphi_1, \varphi_2, \theta) + C_{ji}(\varphi_1, \varphi_2, \theta) + C_{jj}(\varphi_1, \varphi_2, \theta) \\
= \; & \sum_{n=0}^{\infty} \{b_{n,ij+}(\varphi_1, \varphi_2) \cos(n\theta) + a_{n,ij+}(\varphi_1, \varphi_2) \sin(n\theta)\},
\end{aligned} \tag{16}$$

and

$$\begin{aligned}
& C_{ii}(\varphi_1, \varphi_2, \theta) - C_{ij}(\varphi_1, \varphi_2, \theta) - C_{ji}(\varphi_1, \varphi_2, \theta) + C_{jj}(\varphi_1, \varphi_2, \theta) \\
= \; & \sum_{n=0}^{\infty} \{b_{n,ij-}(\varphi_1, \varphi_2) \cos(n\theta) + a_{n,ij-}(\varphi_1, \varphi_2) \sin(n\theta)\},
\end{aligned} \tag{17}$$

where $b_{n,ij+}(\varphi_1, \varphi_2)$ and $b_{n,ij-}(\varphi_1, \varphi_2)$ are covariance functions for each $n \in \mathbb{N}_0$, and $\sum_{n=0}^{\infty} b_{n,ij+}(\varphi_1, \varphi_2)$ and $\sum_{n=0}^{\infty} b_{n,ij-}(\varphi_1, \varphi_2)$ converge. Taking the difference between (16) and (17) and then dividing by 4 yields

$$\frac{C_{ij}(\varphi_1, \varphi_2, \theta) + C_{ji}(\varphi_1, \varphi_2, \theta)}{2} = \sum_{n=0}^{\infty} \{b_{n,ij}(\varphi_1, \varphi_2) \cos(n\theta) + a_{n,ij}(\varphi_1, \varphi_2) \sin(n\theta)\},$$

where

$$b_{n,ij} = \frac{1}{4}(b_{n,ij+} - b_{n,ij-}), \quad a_{n,ij} = \frac{1}{4}(a_{n,ij+} - a_{n,ij-}) n \in \mathbb{N}_0.$$

This establishes (6).

In order to verify that $\mathbf{B}_n(\varphi_1, \varphi_2)$ is a covariance matrix function for each $n \in \mathbb{N}_0$, consider an m-variate stochastic process

$$\mathbf{W}_n(\varphi) = \int_0^{2\pi} \frac{\mathbf{Z}(\varphi, \theta) + (\tilde{\mathbf{Z}}(\varphi, \theta))'}{\sqrt{2}} \cos(n\theta) d\theta, \qquad \varphi \in [0, \pi],$$

where $\{\tilde{\mathbf{Z}}(\mathbf{x}), \mathbf{x} \in \mathbb{S}^2\}$ is an independent copy of $\{\mathbf{Z}(\mathbf{x}), \mathbf{x} \in \mathbb{S}^2\}$. Its covariance matrix function is positively proportional to $\mathbf{B}_n(\varphi_1, \varphi_2)$ as follows,

$$\begin{aligned}
& \operatorname{cov}(\mathbf{W}_n(\varphi_1), \mathbf{W}_n(\varphi_2)) \\
= & \int_0^{2\pi} \int_0^{2\pi} \frac{\operatorname{cov}(\mathbf{Z}(\varphi_1, \theta_1), \mathbf{Z}(\varphi_2, \theta_2)) + \left(\operatorname{cov}(\tilde{\mathbf{Z}}(\varphi_1, \theta_1), \tilde{\mathbf{Z}}(\varphi_2, \theta_2))\right)'}{2} \cos(n\theta_1)\cos(n\theta_2) d\theta_1 d\theta_2 \\
= & \int_0^{2\pi} \int_0^{2\pi} \frac{\mathbf{C}(\varphi_1, \varphi_2, \theta_1 - \theta_2) + (\mathbf{C}(\varphi_1, \varphi_2, \theta_1 - \theta_2))'}{2} \cos(n\theta_1)\cos(n\theta_2) d\theta_1 d\theta_2 \\
= & \sum_{k=0}^{\infty} \mathbf{B}_k(\varphi_1, \varphi_2) \int_0^{2\pi} \int_0^{2\pi} (\cos(k\theta_1)\cos(k\theta_2) + \sin(k\theta_1)\sin(k\theta_2)) \cos(n\theta_1)\cos(n\theta_2) d\theta_1 d\theta_2 \\
& + \sum_{k=0}^{\infty} \mathbf{A}_k(\varphi_1, \varphi_2) \int_0^{2\pi} \int_0^{2\pi} (\sin(k\theta_1)\cos(k\theta_2) - \cos(k\theta_1)\sin(k\theta_2)) \cos(n\theta_1)\cos(n\theta_2) d\theta_1 d\theta_2 \\
= & \begin{cases} 4\pi^2 \mathbf{B}_0(\varphi_1, \varphi_2), & n = 0, \\ \pi^2 \mathbf{B}_n(\varphi_1, \varphi_2), & n \in \mathbb{N}. \end{cases}
\end{aligned}$$

4.4 Proof of Theorem 5

For a fixed $t \in \mathbb{T}$, consider two purely spatial random fields $\{\mathbf{Z}(\mathbf{x}; 0) + \mathbf{Z}(\mathbf{x}; t), \mathbf{x} \in \mathbb{S}^2\}$ and $\{\mathbf{Z}(\mathbf{x}; 0) - \mathbf{Z}(\mathbf{x}; t), \mathbf{x} \in \mathbb{S}^2\}$. Their covariance matrix functions are, in terms of $\mathbf{C}(\varphi_1, \varphi_2, \theta; t)$,

$$\begin{aligned}
& \operatorname{cov}\left(\mathbf{Z}(\varphi_1, \theta_1; 0) + \mathbf{Z}(\varphi_1, \theta_1; t),\ \mathbf{Z}(\varphi_2, \theta_2; 0) + \mathbf{Z}(\varphi_2, \theta_2; t)\right) \\
= & \ 2\mathbf{C}(\varphi_1, \varphi_2, \theta_1 - \theta_2; 0) + \mathbf{C}(\varphi_1, \varphi_2, \theta_1 - \theta_2; t) + \mathbf{C}(\varphi_1, \varphi_2, \theta_1 - \theta_2; -t),
\end{aligned}$$

and

$$\begin{aligned}
& \operatorname{cov}\left(\mathbf{Z}(\varphi_1, \theta_1; 0) - \mathbf{Z}(\varphi_1, \theta_1; t),\ \mathbf{Z}(\varphi_2, \theta_2; 0) - \mathbf{Z}(\varphi_2, \theta_2; t)\right) \\
= & \ 2\mathbf{C}(\varphi_1, \varphi_2, \theta_1 - \theta_2; 0) - \mathbf{C}(\varphi_1, \varphi_2, \theta_1 - \theta_2; t) - \mathbf{C}(\varphi_1, \varphi_2, \theta_1 - \theta_2; -t), \quad (\varphi_k, \theta_k) \in \mathbb{S}^2,\ k = 1, 2.
\end{aligned}$$

It follows from Theorem 3 that

$$\begin{aligned}
&2\mathbf{C}(\varphi_1,\varphi_2,\theta;0)+\mathbf{C}(\varphi_1,\varphi_2,\theta;t)+\mathbf{C}(\varphi_1,\varphi_2,\theta;-t)\\
&+\{2\mathbf{C}(\varphi_1,\varphi_2,\theta;0)+\mathbf{C}(\varphi_1,\varphi_2,\theta;t)+\mathbf{C}(\varphi_1,\varphi_2,\theta;-t)\}'\\
&=\sum_{n=0}^{\infty}\left(\mathbf{B}_{n+}(\varphi_1,\varphi_2;t)\cos(n\theta)+\mathbf{A}_{n+}(\varphi_1,\varphi_2;t)\sin(n\theta)\right),
\end{aligned}\tag{18}$$

and

$$\begin{aligned}
&2\mathbf{C}(\varphi_1,\varphi_2,\theta;0)-\mathbf{C}(\varphi_1,\varphi_2,\theta;t)-\mathbf{C}(\varphi_1,\varphi_2,\theta;-t)\\
&+\{2\mathbf{C}(\varphi_1,\varphi_2,\theta;0)-\mathbf{C}(\varphi_1,\varphi_2,\theta;t)-\mathbf{C}(\varphi_1,\varphi_2,\theta;-t)\}'\\
&=\sum_{n=0}^{\infty}\left(\mathbf{B}_{n-}(\varphi_1,\varphi_2;t)\cos(n\theta)+\mathbf{A}_{n-}(\varphi_1,\varphi_2;t)\sin(n\theta)\right),
\end{aligned}\tag{19}$$

$$(\varphi_k,\theta_k)\in\mathbb{S}^2,\ k=1,2,t\in\mathbb{T}.$$

Taking the difference between (18) and (19) results in (10), with

$$\mathbf{B}_n(\varphi_1,\varphi_2;t)=\frac{1}{4}\mathbf{B}_{n+}(\varphi_1,\varphi_2;t)-\frac{1}{4}\mathbf{B}_{n-}(\varphi_1,\varphi_2;t),$$

and

$$\mathbf{A}_n(\varphi_1,\varphi_2;t)=\frac{1}{4}\mathbf{A}_{n+}(\varphi_1,\varphi_2;t)-\frac{1}{4}\mathbf{A}_{n-}(\varphi_1,\varphi_2,\theta;t),\qquad n\in\mathbb{N}_0.$$

Obviously, $\mathbf{B}_n(\varphi_1,\varphi_2;t)$ is symmetric, and $\sum_{n=0}^{\infty}\mathbf{B}_n(\varphi_1,\varphi_2;t)$ converges.

In order to verify that $\mathbf{B}_n(\varphi_1,\varphi_2;t)$ is a spatio-temporal covariance matrix function on $[0,\pi]\times\mathbb{T}$ for each fixed $n\in\mathbb{N}_0$, consider an m-variate random field

$$\mathbf{W}_n(\varphi;t)=\int_0^{2\pi}\frac{\mathbf{Z}(\varphi,\theta;t)+\mathbf{Z}(\varphi,\theta;-t)+(\tilde{\mathbf{Z}}(\varphi,\theta;t)+\tilde{\mathbf{Z}}(\varphi,\theta;-t))'}{2}\cos(n\theta)d\theta,\quad\varphi\in[0,\pi],\ t\in\mathbb{T},$$

where $\{\tilde{\mathbf{Z}}(\varphi,\theta;t),\varphi\in[0,\pi],\theta\in[0,2\pi],t\in\mathbb{T}\}$ is an independent copy of $\{\mathbf{Z}(\varphi,\theta;t),\varphi\in[0,\pi],\theta\in[0,2\pi],t\in\mathbb{T}\}$. The covariance matrix function of $\{\mathbf{W}_n(\varphi;t),\varphi\in[0,\pi],t\in\mathbb{T}\}$ is a positive scalar product of $\mathbf{B}_n(\varphi_1,\varphi_2;t)$, since

$$\begin{aligned}
&\operatorname{cov}(\mathbf{W}_n(\varphi_1;t_1),\mathbf{W}_n(\varphi_2;t_2))\\
=\ &\frac{1}{2}\int_0^{2\pi}\int_0^{2\pi}\{\operatorname{cov}\left(\mathbf{Z}(\varphi_1,\theta_1;t_1)+\mathbf{Z}(\varphi_1,\theta_1;-t_1),\ \mathbf{Z}(\varphi_2,\theta_2;t_2)+\mathbf{Z}(\varphi_2,\theta_2;-t_2)\right)\\
&+\operatorname{cov}(\tilde{\mathbf{Z}}'(\varphi_1,\theta_1;t_1)+\tilde{\mathbf{Z}}'(\varphi_1,\theta_1;-t_1),\ \tilde{\mathbf{Z}}'(\varphi_2,\theta_2;t_2)+\tilde{\mathbf{Z}}'(\varphi_2,\theta_2;-t_2))\}\cos(n\theta_1)\cos(n\theta_2)d\theta_1d\theta_2\\
=\ &\frac{1}{2}\int_0^{2\pi}\int_0^{2\pi}\{\mathbf{C}(\varphi_1,\varphi_2,\theta_1-\theta_2;t_1-t_2)+\mathbf{C}(\varphi_1,\varphi_2,\theta_1-\theta_2;t_2-t_1)
\end{aligned}$$

$$
\begin{aligned}
&+(\mathbf{C}(\varphi_1,\varphi_2,\theta_1-\theta_2;t_1-t_2)+\mathbf{C}(\varphi_1,\varphi_2,\theta_1-\theta_2;t_2-t_1))'\}\cos(n\theta_1)\cos(n\theta_2)d\theta_1 d\theta_2\\
=&\sum_{n=0}^{\infty}\int_0^{2\pi}\int_0^{2\pi}\{\mathbf{B}_n(\varphi_1,\varphi_2;t)\cos(n(\theta_1-\theta_2))+\mathbf{A}_n(\varphi_1,\varphi_2;t)\sin(n(\theta_1-\theta_2))\}\cos(n\theta_1)\cos(n\theta_2)d\theta_1 d\theta_2\\
=&\begin{cases}4\pi^2\mathbf{B}_0(\varphi_1,\varphi_2;t_1-t_2), & n=0,\\ \pi^2\mathbf{B}_n(\varphi_1,\varphi_2;t_1-t_2), & n\in\mathbb{N}.\end{cases}
\end{aligned}
$$

5 Proof of Theorem 6

It suffices to prove the last part, while the proof of the first part is analogous to that of Theorem 2. If all $\mathbf{V}_{n1}(\varphi;t)$ and $\mathbf{V}_{n2}(\varphi;t)$ ($n\in\mathbb{N}_0$) are normally distributed, then $\mathbf{Z}(\varphi,\theta;t)$ defined by (9) is normally distributed, too. On the other hand, if $\mathbf{Z}(\varphi,\theta;t)$ defined by (9) is normally distributed, then, for each $n\in\mathbb{N}_0$, $\mathbf{Z}(\varphi,\theta;t)\cos(n\theta)$ and $\mathbf{Z}(\varphi,\theta;t)\sin(n\theta)$ are normally distributed as well. So are

$$
\begin{aligned}
&\int_0^{2\pi}\mathbf{Z}(\varphi,\theta;t)\cos(n\theta)d\theta\\
=&\int_0^{2\pi}\sum_{k=0}^{\infty}(\mathbf{V}_{k1}(\varphi;t)\cos(k\theta)+\mathbf{V}_{k2}(\varphi;t)\sin(k\theta))\cos(n\theta)d\theta\\
=&\sum_{k=0}^{\infty}\mathbf{V}_{k1}(\varphi;t)\int_0^{2\pi}\cos(k\theta)\cos(n\theta)d\theta+\sum_{k=0}^{\infty}\mathbf{V}_{k2}(\varphi;t)\int_0^{2\pi}\sin(k\theta)\cos(n\theta)d\theta\\
=&\begin{cases}2\pi\mathbf{V}_{01}(\varphi;t), & n=0,\\ \pi\mathbf{V}_{n1}(\varphi;t), & n\in\mathbb{N},\end{cases}
\end{aligned}
$$

and

$$
\begin{aligned}
&\int_0^{2\pi}\mathbf{Z}(\varphi,\theta;t)\sin(n\theta)d\theta\\
=&\sum_{k=0}^{\infty}\mathbf{V}_{k1}(\varphi;t)\int_0^{2\pi}\cos(k\theta)\sin(n\theta)d\theta+\sum_{k=0}^{\infty}\mathbf{V}_{k2}(\varphi;t)\int_0^{2\pi}\sin(k\theta)\sin(n\theta)d\theta\\
=&\begin{cases}0, & n=0,\\ \pi\mathbf{V}_{n2}(\varphi;t), & n\in\mathbb{N}.\end{cases}
\end{aligned}
$$

5.1 Proof of Theorem 7

According to Theorem 8 of [24], it suffices to check inequality (1) for (15). For every $l \in \mathbb{N}$, any $\mathbf{w}_i \in \mathbb{R}^m$, $\varphi_i \in [0, \pi]$, $\theta_i \in [0, 2\pi]$, and $t_i \in \mathbb{T}$ $(k = 1, \ldots, l)$, we have

$$
\begin{aligned}
& \sum_{i=1}^{l}\sum_{j=1}^{l} \mathbf{w}_i' \mathbf{C}(\varphi_i, \varphi_j, \theta_i - \theta_j; t_i - t_j)\mathbf{w}_j \\
= & \sum_{n=0}^{\infty}\sum_{i=1}^{l}\sum_{j=1}^{l} \mathbf{w}_i' (\mathbf{B}_n(\varphi_i, \varphi_j; t_i - t_j)\cos(n(\theta_i - \theta_j)) + \mathbf{A}_n(\varphi_i, \varphi_j; t_i - t_j)\cos(n(\theta_i - \theta_j)))\mathbf{w}_j \\
= & \sum_{n=0}^{\infty}\sum_{i=1}^{l}\sum_{j=1}^{l} \big\{ \mathbf{w}_i' \cos(n\theta_i)\mathbf{B}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{w}_j \cos(n\theta_j) + \mathbf{w}_i' \sin(n\theta_i)\mathbf{B}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{w}_j \sin(n\theta_j) \\
& + \mathbf{w}_i' \sin(n\theta_i)\mathbf{A}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{w}_j \cos(n\theta_j) - \mathbf{w}_i' \cos(n\theta_i)\mathbf{A}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{w}_j \sin(n\theta_j) \big\} \\
= & \sum_{n=0}^{\infty}\sum_{i=1}^{l}\sum_{j=1}^{l} (\mathbf{u}_{ni}' \mathbf{B}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{u}_{nj} + \mathbf{v}_{ni}' \mathbf{B}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{v}_{nj} \\
& + \mathbf{v}_{ni}' \mathbf{A}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{u}_{nj} - \mathbf{u}_{ni}' \mathbf{A}_n(\varphi_i, \varphi_j; t_i - t_j)\mathbf{v}_{nj}) \\
\geq & \ 0,
\end{aligned}
$$

where the last inequality follows from the assumption (14), $\mathbf{u}_{ni} = \mathbf{w}_i \cos(n\theta_i), \mathbf{v}_{ni} = \mathbf{w}_i \sin(n\theta_j)$.

References

[1] Andrews, G. E., Askey, R., Roy, R.: Special Functions. Cambridge University Press, Cambridge (1999)

[2] Askey, R., Bingham, N. H.: Gaussian processes on compact symmetric spaces. Z. Wahrscheinlichkeitstheorie verw. Gebiete **37**, 127-143 (1976)

[3] Bingham, N. H.: Positive definite functions on spheres. Proc. Cambridge Phil. Soc. **73**, 145-156 (1973)

[4] Castruccio, S. and Stein, M. L.: Global space-time models for climate ensembles. Ann. Appl. Stat. **7**, 1593-1611 (2013)

[5] Cheng, D., Xiao, Y.: Excursion probability of Gaussian random fields on sphere. Bernoulli **22**, 1113-1130 (2016)

[6] Cohen, S., Lifshits, M. A.: Stationary Gaussian random fields on hyperbolic spaces and on Euclidean spheres. ESAIM **16**, 165-221 (2012)

[7] D'Ovidio, M.: Coordinates changed random fields on the sphere. J. Stat. Phys. **154**, 1153-1176 (2014)

[8] Du, J., Ma, C., Li, Y.: Isotropic variogram matrix functions on spheres. Math. Geosci. **45**, 341-357 (2013)

[9] Gangolli, R.: Positive definite kernels on homogeneous spaces and certain stochastic processes related to Lévy's Brownian motion of several parameters. Ann Inst H Poincaré B **3**, 121-226 (1967)

[10] Gaspari, G., Cohn, S. E.: Construction of correlations in two and three dimensions. Q. J. R. Meteorol. Soc. **125**, 723-757 (1999)

[11] Gaspari, G., Cohn, S. E., Guo, J., Pawson, S.: Construction and application of covariance functions with variable length-fields. Q. J. R. Meteorol. Soc. **132**, 815-1838 (2006)

[12] Gradshteyn, I. S., Ryzhik, I. M.: Tables of Integrals, Series, and Products, 7th edtion. Academic Press, Amsterdam (2007)

[13] Hannan, E. J.: Multiple Time Series. Wiley, New York (1970)

[14] Hitczenko, M. and Stein, M. L.: Some theory for anisotropic processes on the sphere. Statist. Meth. **9**, 211-227 (2012)

[15] Huang, C., Zhang, H., and Robeson, S.: A simplified representation of the covariance structure of axially symmetric processes on the sphere. Statist. Probab. Lett. **82**, 1346-1351 (2012)

[16] Jones, R. H.: Stochastic processes on a sphere. Ann. Math. Statist. **34**, 213-218 (1963)

[17] Jun, M.: Non-stationary cross-covariance models for multivariate processes on a globe. Scand. J. Statist., **38**, 726-747 (2011)

[18] Jun, M. and Stein, M. L.: An approach to producing space-time covariance functions on spheres. Technometrics **49**, 468-479 (2007)

[19] Jun, M. and Stein, M. L.: Nonstationary covariance models for global data. Ann. Appl. Statist. **2**, 1271-1289 (2008)

[20] Lamberg, L., Muinonen, K., Ylönen, J., Lumme, K.: Spectral estimation of Gaussian random circles and spheres. J. Comput. Appl. Math. **136**, 109-121 (2001)

[21] Leonenko, N., Sakhno, L.: On spectral representation of tensor random fields on the sphere. Stoch. Anal. Appl. **31**, 167-182 (2012)

[22] Leonenko, N., Shieh, N.: Rényi function for multifractal random fields. Fractals, **21**, 1350009, 13 pp (2013)

[23] Ma, C.: Vector random fields with second-order moments or second-order increments. Stoch. Anal. Appl. **29**, 197-215 (2011)

[24] Ma, C.: Stationary and isotropic vector random fields on spheres. Math. Geosci. **44**, 765-778 (2012)

[25] Ma, C.: Isotropic covariance matrix functions on all spheres. Math. Geosci. **47**, 699-717 (2015)

[26] Ma, C.: Stochastic representations of isotropic vector random fields on spheres. Stoch. Anal. Appl. **34**, 389-403 (2016)

[27] Ma, C.: Isotropic covariance matrix polynomials on spheres. Stoch. Anal. Appl., To appear.

[28] Ma, C.: Time varying isotropic vector random fields on spheres. J. Theor. Probab., To appear (DOI 10.1007/s10959-016-0689-1).

[29] Malyarenko, A.: Invariant Random Fields on Spaces with a Group Action. Springer, New York (2013)

[30] Malyarenko, A., Olenko, A.: Multidimensional covariant random fields on commutative locally compact groups. Ukrainian Math. J. **44**, 1384-1389 (1992)

[31] McLeod, M. G.: Stochastic processes on a sphere. Phy. Earth Plan. Interior **43**, 283-299 (1986)

[32] Mokljacuk, M. P., Jadrenko, M. I.: Linear statistical problems for stationary isotropic random fields on a sphere, I. Theor. Prob. Math. Statist. **18**, 115-124 (1979)

[33] Roy, R.: Spetral analysis for random process on the circle. J. Appl. Prob. **9**, 745-757 (1972)

[34] Roy, R.: Spectral analysis for a random process on the sphere. Ann. Inst. Statist. Math. **28**, 91-97 (1976)

[35] Schoenberg, I.: Positive definite functions on spheres. Duke Math. J. **9**, 96-108 (1942)

[36] Stein, M.: Spatial variation of total column ozone on a global scale. Ann. Appl. Statist. **1**, 191-210 (2007)

[37] Szegö, G. Orthogonal Polynomials, 4th edition. Amer. Math. Soc. Colloq. Publ., vol 23. Amer. Math. Soc., Providence (1975)

[38] Watson, G. N.: A Treatise on the Theory of Bessel Functions, 2nd edn. Cambridge University Press, London (1944)

[39] Yadrenko, A. M. Spectral Theory of Random Fields. Optimization Software, New York (1983)

[40] Yaglom, A. M.: Second-order homogeneous random fields. Proc. 4th Berkeley Symp. Math. Stat. Prob. **2**, 593-622 (1961)

[41] Yaglom, A. M.: Correlation Theory of Stationary and Related Random Functions. vol. I. Springer, New York (1987)